

Project Title

Personal Protective Equipment Pictorial Guide

Project Lead and Members

- Lim Hui Yun
- Foo Cui Shan
- Yuen Wyn Yuen

Organisation(s) Involved

Singapore National Eye Centre

Healthcare Family Group(s) Involved in this Project

Nursing

Applicable Specialty or Discipline

Infection Control

Project Period

Start date: January 2023

Completed date: April 2023

Aim(s)

- To create a resource that health care workers (HCWs) can use to easily refer to and apply correct PPE usage for different infectious diseases.
- To enhance HCWs' knowledge in applying correct isolation precautions to prevent HAIs.
- To improve workplace safety and patient safety by promoting proper PPE use.

Background

See poster appended/ below

Methods

See poster appended/ below

Results

See poster appended/ below

Conclusion

See poster appended/ below

Project Category

Care & Process Redesign

Risk Management, Preventive Approach

Keywords

Personal Protective Equipment (PPE), Pictorial Guide, Healthcare Workers

Name and Email of Project Contact Person(s)

Name: Lim Hui Yun

Email: singaporehealthcaremanagement@singhealth.com.sg



Personal Protective Equipment Pictorial Guide


Singapore Healthcare Management 2023

Lim Hui Yun ¹, Foo Cui Shan ², Yuen Wyn Yuen ³
 1. Singapore National Eye Centre, Outpatient Clinic
 2. Singapore National Eye Centre, Infection Control Nurse
 3. Singapore National Eye Centre, Operating Theatre

Introduction

Personal protective equipment (PPE) is crucial in preventing healthcare-associated infection (HAIs) and ensuring workplace safety.¹ However, proper use of PPE can be challenging, especially when dealing with different infectious diseases for accurate and appropriate use of PPE with reference to the lengthy policies and guidelines. Therefore the use of a pictorial guide was designed to provide a guidance for reference for Health Care Workers (HCWs) in SNEC.

Ishikawa Cause and Effect Diagram



Suárez-Barraza & Rodríguez-González, 2018.²

Objectives

- To create a resource that health care workers (HCWs) can use to easily refer to and apply correct PPE usage for different infectious diseases.
- To enhance HCWs' knowledge in applying correct isolation precautions to prevent HAIs.
- To improve workplace safety and patient safety by promoting proper PPE use.


Methodology

The study was conducted in SNEC with 83 HCWs between January to April in year 2023 from different healthcare professions inclusive of allied health professionals, ancillary and clinical staffs.

The PPE pictorial guide included images of different types of PPE and their correct usage for various infectious disease, such as COVID-19, Tuberculosis (TB), and Methicillin-resistant Staphylococcus aureus (MRSA).

Pre and post implementation surveys were conducted to assess the effectiveness of the intervention. The feedback from HCWs was also collected to evaluate the acceptability of the PPE pictorial guide

PPE Pictorial Guide Pre and Post Implementation Survey



Results


The result showed a significant improvement in HCWs' knowledge in applying correct isolation precautions after the implementation of the PPE pictorial guide in comparison to guideline from institutional policies. **The mean score for knowledge increased from 54% in the pre-implementation survey to 81% in the post-implementation survey.** Feedback from HCWs was also positive, with all participants finding the PPE pictorial guide easy to understand and use indicating the effectiveness of the intervention in improving workplace safety and preventing HAIs.

PPE Pictorial Guide

Disease-specific isolation recommendations			Surgical mask	N95 Mask / Powered air purifier respirator	Isolation gown with cuff	Gloves	Eye protector
Airborne precaution	Monkey Pox	Duration of illness		✓	✓	✓	✓
	Pulmonary Tuberculosis <i>Active and suspect</i>	Discontinue precaution only when patient on effective therapy for 14 days and 3 consecutive sputum smears negative for acid-fast bacilli		✓	✓	✓	✓
	Measles	4 days after onset of rash; duration of illness in immune compromised		✓	✓	✓	
Droplet precaution	Mumps / Rubella	Until 5 days after the onset of swelling		✓	✓	✓	
Airborne + Contact precaution	Herpes Zoster (Varicella Zoster/ Shingles) <i>Duration of illness</i>	Duration of illness		✓	✓	✓	✓
	Coronavirus associated with severe acute respiratory syndrome (SARS-CoV)	From onset of symptoms		✓	✓	✓	✓
Contact Precaution	Clostridium difficile	Duration of illness + 2 days after last unformed stool	✓		✓	✓	
	Conjunctivitis	Bacterial pink eye: after 24 hours of antibiotic treatment Viral pink eye: 1 week	✓		✓	✓	
	Human Immunodeficiency syndrome (HIV)	Duration of illness	✓		✓	✓	
	Hepatitis B	Duration of illness	✓		✓	✓	
	Multidrug-resistant organisms (MDROs) e.g. MRSA, VRE, CP-CRE	Duration of illness	✓		✓	✓	
	Herpes Zoster (Varicella Zoster/ Shingles) <i>Until lesions dry and crusted</i>	Until lesions dry and crusted	✓		✓	✓	
				✓		✓	

Eye protector are to be worn if there is possibility of splashes, sprays, respiratory droplet exposure and during aerosol generating procedure (AGP)
<https://www.cdc.gov/infectioncontrol/guidelines/isolation/appendix/type-duration-precautions.html#T>
 Brought to you by: SNEC Infection Control Committee

Continuous PDSA Cycle



Repeat the PDSA cycle if further refinements are required, incorporating the revised guide and assessing its impact on HCWs' knowledge and confidence. By following the PDSA cycle³, the project can continuously assess, refine, and optimize the PPE pictorial guide to effectively enhance HCWs' knowledge and confidence in applying correct isolation precautions. This iterative approach ensures that the guide remains relevant, accessible, and impactful, ultimately improving workplace safety and preventing HAIs in the healthcare setting.

Conclusion

The PPE pictorial guide was an effective intervention in enhancing HCW's knowledge and confidence in applying correct isolation precautions in preventing HAIs. The easy-to-refer format of the guide made it accessible and usable for HCWs from different professions. The positive feedback from HCWs and the significant improvement in their knowledge and confidence underscore the importance of providing such resources to improve workplace safety and patient safety. The findings of this study have important implications for healthcare facilities seeking to enhance their infection prevention and control measures.

References
 1. Honda H, Iwata K: Personal protective equipment and improving compliance among healthcare workers in high-risk settings. *Curr Opin Infect Dis.* 2016, 29:400-6. [10.1097/QCO.0000000000000280](https://doi.org/10.1097/QCO.0000000000000280)
 2. Suárez-Barraza, M. F., & Rodríguez-González, F. G. (2018). Cornerstone root causes through the analysis of the Ishikawa diagram, is it possible to find them? *International Journal of Quality and Service Sciences*, 11(2), 302–316. <https://doi.org/10.1108/ijqss-12-2017-0113>
 3. Taylor, M. J., McNicholas, C., Nicolay, C., Darzi, A., Bell, D., & Reed, J. E. (2013). Systematic Review of the Application of the Plan-Do-Study-Act Method to Improve Quality in Healthcare. *BMJ Quality & Safety*, 23(4), 290–298. <https://doi.org/10.1136/bmjqs-2013-001862>