HEALTHCARE INNOVATION

CHI Learning & Development (CHILD) System

Project Title

Initiatives to Reduce Blood Specimen Rejection Rate

Project Lead and Members

Project lead: Lim Jia Ying

Project members: Samantha Koh, Cuevas Sheryll Alojado, Lew Shu Ling, Tacbianan

Grace Rafanan, Sa'adiah Binte Arsad, Henry Teo, Lyon Loo

Organisation(s) Involved

Ng Teng Fong General Hospital

Healthcare Family Group Involved in this Project

Nursing, Healthcare Administration

Applicable Specialty or Discipline

Medical & Laboratory Technology

Aims

Ward B16S intends to reduce the median blood specimen rejection rate from 0.39% to 0.30% in Ward B16s by May 2020 for Ward B16S patients because we want to prevent delay in treatments and to reduce patients' complaints.

Background

See poster appended/below

Methods

See poster appended/below

Results

See poster appended/ below



CHI Learning & Development (CHILD) System

Lessons Learnt

The creation of visual cues and reinforcements on the correct sequences of venipuncture has ensured the quality of the specimens.

Conclusion

The compliance to the correct sequence of draw for blood tubes and the standardisation of 8 inversions for all blood tubes had shown positive results in reducing blood specimens rejection rates significantly

Project Category

Care & Process Redesign, Quality Improvement, Job Effectiveness

Keywords

Blood Specimen Rejection, Inpatient Wards

Name and Email of Project Contact Person(s)

Name: Lim Jia Ying

Email: jeanette limjy@hotmail.com

[Restricted, Non-sensitive]

INITIATIVES TO REDUCE BLOOD SPECIMEN REJECTION RATE

LIM JIA YING, SAMANTHA KOH CHU PING, CUEVAS SHERYLL ALOJADO, LEW SHU LING, TACBIANAN GRACE RAFANAN, SA'ADIAH BINTE ARSAD, HENRY TEO WEE MENG, LYON LOO JUN YUAN

Ward B16 Subsidised, Ng Teng Fong General Hospital

SAFETY QUALITY PATIENT EXPERIENCE



Define Problem, Set Aim

Problem statement

Between January 2018 to April 2019, Ward B16s saw an increasing trend of specimen rejection. Ranking Ward B16s as the top 6 rejections in Ng Teng Fong General Hospital with the highest blood specimen rejections on several occasions starting from October 2018 onwards. The median shift from 0.27% (April 2018 – September 2018) to 0.39% (October 2018 - April 2019). The median rejection across NTFGH is about 0.35%.

This can lead to the following:


- 1. Delay in treatment
- 2. Patient complaints (caused by undue pain)
- 3. Reworking and increase waste

Aim


Ward B16s intends to reduce the median blood specimen rejection rate from 0.39% to 0.30% in Ward B16s by May 2020 for Ward B16s patients because we want to prevent delay in treatments and to reduce patients' complaints.

Outcome measure: (total number of blood rejected in Ward B16s (monthly) (total number of rejected specimens in NTFGH (monthly)

Establish Measures




Analyse Problem



What are the probable <u>root</u> causes?


Fishbone diagram and 5-Why were used to derive the potential root causes. These key root causes are identified using pareto chart shown below.



Ng Teng Fong General Hospital



Select Changes





Test & Implement Changes

How do we pilot the changes? What are the initial results?

CYCLE	PLAN	DO	STUDY	ACT
1	NIC to do visual check for all specimens prior to dispatch.	Carried out from 10 December 2018 till 14 March 2019.	The blood specimen rejection rates decreased in January 2019.	Adapt
2	Develop visual cues for users doing venepuncture	Carried out from 9 July 2019.	Consecutive 6 months of achieving targeted rejection rates of below 0.3% from December 2019 till May 2020.	Adopt

To ensure specimens quality and to provide reinforcement on the venepuncture process, the team has developed visual cues targeting on the correct sequence of draw for blood tubes and has standardised the number of inversions to 8 inversions for all blood tubes. Visual cues were attached on the phlebotomy trolley laptop.





Conclusion

Audit was conducted in Ward B16s from December 2019 – May 2020.

The compliance to the correct sequence of draw for blood tubes and the standardisation of 8 inversions for all blood tubes had shown positive results in reducing blood specimens rejection rates significantly.

Ward B16s has achieved the targeted blood specimen rejection rate of less than 0.30% for 6 consecutive months.